

Impact of Hard Palate Angulation Caused by Septal Deviation on Maxillary Sinus Volume

Sert Damakta Açılanmaya Neden Olan Septum Deviasyonunun Maksiller Sinüs Hacmine Etkisi

Emrah Sapmaz¹, Ahmet Kavaklı², Hilal Irmak Sapmaz³, Murat Ögetürk²

¹Department of Otorhinolaryngology, Gaziosmanpaşa University School of Medicine, Tokat, Turkey

²Department of Anatomy, Firat University School of Medicine, Elazığ, Turkey

³Department of Anatomy, Gaziosmanpaşa University School of Medicine, Tokat, Turkey

Original Investigation

Özgün Araştırma

Abstract

Objective: To investigate the effect of hard palate angulation caused by septal deviation on the volume of the maxillary sinus.

Methods: Coronal computed tomographic (CT) scans of 1568 patients aged from 18 to 60 were examined. CT scans of 402 patients were included in the study. On these scans, the maxillary sinus volume, the angle of the nasal septal deviation, and the angulation of the hard palate were calculated using the ImageJ software. Each maxillary sinus volume was statistically compared with each other and with those in the control group. Correlations between palatal angulation and septal deviation were determined.

Results: Deviated nasal septum whether with or without deflection of the hard palate was noted to have caused changes in the volume of the maxillary

sinus in both female and male patients. The volume of the maxillary sinus on the deviated side was less than that of the opposite side, and the differences between the volumes of both sinuses were statistically significant ($p < 0.05$). No significant differences were noted when compared with the control group. A positive correlation was observed between the nasal septal deviation angle and the angulation of the hard palate.

Conclusion: Regardless of whether or not it affects the hard palate, nasal septal deviation reduces the volume of the maxillary sinus on the deviated side but does not affect the total volume of the maxillary sinuses. Significant differences between the volumes on the two sides can lead to facial asymmetry.

Keywords: Nasal septal deviation, hard palate, maxillary sinus, computed tomography



ORCID IDs of the authors:

E.S. 0000-0002-0346-8704;
A.K. 0000-0003-0754-8901;
H.I.S. 0000-0002-9821-1628.

Öz

Cite this article as: Sapmaz E, Kavaklı A, Sapmaz H, Ögetürk M. Impact of Hard Palate Angulation Caused by Septal Deviation on Maxillary Sinus Volume. Turk Arch Otorhinolaryngol 2018; 56(2): 75-80.

This study was presented at the 24th International Symposium on Morphological Sciences (ISMS 2015), September 2-9, 2015, Istanbul, Turkey.

Bu çalışma, 24. Uluslararası Morfolojik Bilimler Sempozyumunda sunulmuştur, 2-9 Eylül 2015, İstanbul, Türkiye.

Corresponding Author/Sorumlu Yazar: Emrah Sapmaz; emrhils@yahoo.com

Received Date/Geliş Tarihi: 25.10.2017

Accepted Date/Kabul Tarihi: 16.03.2018

© Copyright 2018 by Official Journal of the Turkish Society of Otorhinolaryngology and Head and Neck Surgery Available online at www.turkarchotolaryngol.net

© Telif Hakkı 2018 Türk Kulak Burun Boğaz ve Baş Boyun Cerrahisi Derneği Makale metnine www.turkarchotolaryngol.net web sayfasından ulaşılabilir.

DOI: 10.5152/tao.2018.2987

Amaç: Sert damakta açılanmaya neden olan septum deviasyonunun maksiller sinüs hacmine olan etkisini araştırmak.

Yöntemler: Yaşları 18 ile 60 arasında değişen 1568 hastanın koronal planda çekilen paranasal sinüs tomografileri incelendi. Toplam 402 hasta tomografisi çalışmaya dâhil edildi. Tomografi görüntülerinden ImageJ programı kullanılarak maksiller sinüs hacimleri, septum deviasyonu ve sert damak açıları ölçüldü. Maksiller sinüs hacimleri kendi aralarında ve kontrol grubu ile istatistiksel olarak karşılaştırıldı. Sert damak ve septal açılanmalar arasındaki korelasyon araştırıldı.

Bulgular: Hem erkek hem kadın hasta grubunda sert damakta açılanmaya yol açan veya açmayan septum deviasyonu kişilerin maksiller sinüs hacimlerinde farklılıklar oluşturmuştu. Deviasyonun olduğu tarafta-

ki maksiller sinüs hacmi deviasyonun karşı tarafındaki maksiller sinüs hacminden daha küçük olarak ölçüldü ve bu fark istatistiksel olarak anlamlı idi ($p < 0.05$), fakat kontrol grubu ile karşılaştırıldıklarında aralarında istatistiksel anlamlı bir fark tespit edilmedi. Septum deviasyonu açısı ile sert damakta meydana gelen açılanma arasında pozitif bir korelasyon tespit edildi.

Sonuç: Septum deviasyonu sert damağı etkilesin veya etkilemesin, deviasyon tarafındaki maksiller sinüs hacmini küçültmekte fakat total maksiller sinüs hacmini etkilememektedir. Bu farkın çok fazla olduğu kişilerde fasiyal asimetri dahi tespit etmek mümkün olabilir.

Anahtar kelimeler: Septum deviasyonu, sert damak, maksiller sinüs, bilgisayarlı tomografi

Introduction

The development of the paranasal sinuses is yet to be elucidated. It is argued that nasal airflow, brain development, muscle strength, and migration play significant roles in their formation (1, 2). Volumes

of the maxillary sinus and its anatomical neighborhood depend on aeration (3). Maxillary sinus development is directly associated with the alveolar process and hard palate. Changes in volume after full development are associated with chronological

and pathological conditions (4). Decreased airflow through the nasopharynx reduces oxygen pressure and negatively affects paranasal sinus development (5).

Many deformities of the nasal septum are considered as developmental defects. These defects are classified as nasal septal deviations. Babyhood and childhood traumas that are often considered insignificant and go unnoticed are suspected to be the likely causes of several developmental deformities of the nasal septum. Trauma in early life can lead to asymmetry in the entire nasal structure depending on the degree of bending and deviation of the nasal septal cartilage (6).

In previous studies, nasal septal deviation has been shown to affect the volume of the maxillary sinus (7, 8), but to the best of our knowledge, the palatine bone has not yet been evaluated. A good understanding of the developmental variations of the paranasal sinuses will enable us to better comprehend the period of diseases and provide information in the decision-making process for a surgical intervention and the type of surgical procedure (9).

In this study, we calculated the volumes of maxillary sinuses on computed tomographic (CT) scans of paranasal sinuses using the Cavalieri's principle to explore whether nasal septal deviations that deflect the hard palate affect the volume of the maxillary sinus.

Methods

From January 2012 to December 2013, 1568 patients aged from 18 to 60 years consulted our otorhinolaryngology clinic with symptoms of headache and inability to breathe through the nose. After obtaining approval from the Research Ethics Committee (2014-06/10) and the department of Radiology, CT scans of the patients' paranasal sinuses taken in the coronal plane were

examined. Scans of 1166 patients who presented with damaged anatomical structure caused by previous sinonasal surgery, nasal polyposis, sinonasal tumors or pansinusitis, or whose scans were not properly performed were excluded from the study. We did not obtain informed consent from the patients since this is a retrospective study. Ultimately, CT scans of 202 male and 200 female patients were included in the study.

Patients were assigned to five separate groups as follows:

Group A (n=79): Patients with a right septal deviation accompanied by hard palate angulation.

Group B (n=80): Patients with a left septal deviation accompanied by hard palate angulation.

Group C (n=83): Patients with only right septal deviation.

Group D (n=79): Patients with only left septal deviation.

Group E (n=81): Control patients with no sinonasal pathology.

The evaluation of CT scans were performed in the coronal plane of 5-mm slices (Somatom X; Siemens, Erlangen, Germany) (scan settings: 110 kV, 200 mA, scan field of view [FoV]: 170 mm). Maxillary sinus volumes, nasal septal deviation, and deviation of the hard palate were calculated using ImageJ software (ImageJ, 1.49v: <http://rsb.info.nih.gov/ij/>) (Figure 1).

Measuring the Hard Palate Angle

To eliminate possible imaging errors when measuring hard palate angulation, a line was drawn in the same section between the lesser wings of the sphenoid bone participating in the structure of the orbit on both sides (Figure 2A). This line was projected parallelly so that its segment at the hard palate level (Figure 2B) formed one ray of the angle and the line tangent to the base of the hard palate (Figure 2C) formed another ray of the angle (Figure 2D). Since the lesser wings of the sphenoid bone that




Figure 1. ImageJ program




Figure 2. Measuring the angle of the hard palate




Figure 3. Calculating the volume of the maxillary sinus




Figure 4. Measuring the angle of nasal septal deviation

Table 1. Statistical data of male patients

Groups		Volume of Right Maxillary Sinus (cm ³)(Min-Max)	Volume of Left Maxillary Sinus (cm ³) (Min-Max)	Total Volume of Maxillary Sinus (cm ³)(Min-Max)	Angle of Nasal Septal Deviation (Mean±SD)	Angle of Hard Palate (Mean±SD)	Correlation between Angle of Nasal Septal Deviation and Hard Palate "r"
Group A	(n=40)	9.32	11.39	20.71	14.42±4.73	7.75±2.65	0.504
		(4.47-17.61) ^{a,b}	(5.75-18.22) ^a	(10.22-35.83) ^a			
Group B	(n=39)	11.73	9.40	21.14	14.32±3.28	6.96±1.67	0.328
		(5.04-20.07) ^a	(3.48-19.29) ^{a,b}	(8.52-39.36) ^a			
Group C	(n=41)	9.43	10.98	20.41	13.96±3.75		
		(3.07-13.68) ^{a,b}	(4.18-14.28) ^a	(7.25-27.88) ^a			
Group D	(n=41)	11.28	9.58	20.86	14.12±3.56		
		(4.88-17.37) ^a	(4.58-16.99) ^{a,b}	(9.58-34.36) ^a			
Control (n=41)		10.64	10.69	21.34			
		(3.77-17.01)	(3.23-17.65)	(7.00- 34.03)			

^a: Statistically insignificant compared to the maxillary sinus volumes in the control group (p>0.05)

^b: Statistically significant compared to the volume of the maxillary sinus on the opposite side with no septal deviation (p<0.05)

participate in the structures of the orbits on both sides were taken as a reference, the patient's head was deemed to be improperly positioned and the scan was excluded if both eyeballs did not simultaneously appear on the same slice. Hard palate angulation was measured at the level where the angle of septum deviation was the largest, using the ImageJ software.

Calculating the Maxillary Sinus Volume

Cavalieri's principle was used for calculating the total volumes. Volumes of both maxillary sinuses were calculated using the planimetry method with ImageJ software. Maxillary sinus areas were measured one by one on each slice from the moment

they appeared on the slices. The total number multiplied by the interslice distance equaled the total volume (Figure 3). Data for maxillary sinus volumes were individually compared (with both the deviated side and the other side) and with those of the control group.

Measuring the Nasal Septal Deviation Angle

The curvature of the septum was measured as the angle (Figure 4D) formed by the line drawn between the crista galli (Figure 4A) and the crista nasalis of the maxilla (Figure 4B), and the line drawn through the maximum deviated point (Figure 4C) using ImageJ software.

Table 2. Statistical data of female patients

Groups		Volume of Right Maxillary Sinus (cm3) (Min-Max)	Volume of Left Maxillary Sinus (cm3) (Min-Max)	Total Volume of Maxillary Sinus (cm3) (Min-Max)	Angle of Nasal Septal Deviation (Mean±SD)	Angle of Hard Palate (Mean±SD)	Correlation between Angle of Nasal Septal Deviation and Hard Palate "r"
Group A	(n=39)	7.40	9.00	16.41	13.28±2.61	9.60±3.03	0.684
		(2.43-12.56) ^{a,b}	(3.77-14.87) ^a	(6.33-26.61) ^a			
Group B	(n=41)	9.12	7.57	16.69	13.35±3.10	10.61±2.24	0.199
		(3.55-16.25) ^a	(2.93-12.96) ^{a,b}	(6.48-27.22) ^a			
Group C	(n=42)	7.86	8.89	16.84	12.75±3.76		
		(3.02-15.56) ^{a,b}	(5.46-16.85) ^a	(8.99-32.41) ^a			
Group D	(n=38)	8.80	7.53	16.33	12.92±4.12		
		(4.63-13.22) ^a	(4.07-12.58) ^{a,b}	(8.70-25.05) ^a			
Control (n=40)		8.17	8.51	16.68			
		(3.68-15.34)	(3.53-15.61)	(7.21-30.95)			

^a: Statistically insignificant compared to the maxillary sinus volumes in the control group was ($p>0.05$)

^b: Statistically significant compared to the volume of the maxillary sinus on the opposite side with no septal deviation ($p<0.05$)

Statistical analysis

Data were analyzed using SPSS software. Since the data did not meet the assumptions of parametric tests, groups were compared using the non-parametric Mann-Whitney U test. Correlation between the groups was analyzed with Pearson's correlation. A value of $p<0.05$ was considered statistically significant.

Results

In both the male and female patient groups, nasal septal deviation (right or left) with or without angulation of the os palatinum had caused reduction in the maxillary sinus volume on the deviated side compared with that on the opposite side ($p<0.05$). However, no statistically significant differences were found between the volumes of the deviated and the opposite sides compared with the control group ($p>0.05$). In addition, no statistically significant differences were found between the maxillary sinus volumes of each patient group compared with those of the control group ($p>0.05$). In both the female and the male patient groups, a positive correlation was noted between the angulation of the palatine process and the angle of the nasal septal deviation. Angulation in the palatine process was observed to have increased with nasal septal deviation. Demographic and statistical data of the groups are provided in Tables 1 (male patients) and 2 (female patients).

Discussion

Nasal septal deviations are deformities that present with a deflection, angulation, or luxation of the bones and the cartilage, which form the septal roof. While such deformities are often deemed to be an outcome of nasal micro-fractures, they are also believed to occur as a result of minor facial traumas as well as during the neonatal period (10, 11). Changes in the intrauterine position of the fetus and the newborn, increased transnatal pressure, and traumas during labor are believed to lead to changes in septal development and hence to deviation (10, 12).

The nasal septum has a direct role in the development of the premaxilla and an indirect role in the development of the maxilla (12). Holton et al. (13) showed that septal deviations are associated with the hard palate and the lateral wall asymmetry of the nasal cavity. A study conducted with twins with different septal structures reports that septal deviations affect the antero-posterior development of the nose and the maxilla (14). Nasal septal deviations that arise in the fetal period are also reported to lead to both facial asymmetry and malocclusion (15). While the prevalence of nasal septal deviations is reported to be 20%-31%, a study on patients consulting an otorhinolaryngology clinic for any reason reports this rate as 89.2% (16).

While nasal airflow plays a crucial role in the development of the paranasal sinuses and the craniofacial skeleton (17), positive airflow in the nasopharynx plays an important role in the development of paranasal sinuses. The obstruction of airflow and reduced oxygen pressure interrupt the development of the paranasal sinus (18). Hypertrophy of the pharyngeal tonsil, which causes obstruction in the posterior paranasal sinus, and nasal septal deviation affecting the development of the maxilla can disrupt the development of paranasal sinuses (4). Oral breathing in the absence of nasal airflow and pressing down the mandible and pulling the tongue down and forward affects the development of the maxillofacial skeleton (19).

The development patterns of paranasal sinuses vary by gender and age. While maxillary sinuses can develop differently on the two sides, no significant differences were reported when maxillary sinus volumes were compared between the genders and the volumes of each side (3, 20). However, Karakas and Kavakli (21) and Uchida et al. (22) reported that paranasal sinus volumes differed between genders. Age and alveolar process height are reported to be the major factors affecting the maxillary sinus volume (23). Barghouth et al. (24) report that the right max-

illary sinus is considerably longer than the left maxillary sinus in babies younger than 9 months, and the left maxillary sinus is longer than the right one in children older than eight years. Since the incomplete development of the maxillary sinuses in childhood can be misleading, we included patients older than 18 years in our study.

Factors leading to reduced nasal airflow can also lead to differences in the volumes of paranasal sinuses. Firat et al. (25) report to have found the total volume of ethmoid cells on the deviated side of a nasal septum to be significantly reduced compared with the other side. In our study, we likewise found that maxillary sinus volume on the deviated side of the septum was reduced regardless of an angulation of the hard palate ($p < 0.05$). This suggests that the angulation of the hard palate is caused by the deviated nasal septum rather than the reduced volume of the maxillary sinus, since the presence or absence of an angulation of the hard palate caused no significant differences in the volume of the maxillary sinus. The positive correlation between the angle of the deviation and the angle of the hard palate seems to support our observation. In addition, no difference was reportedly found compared with the control group in the maxillary sinus volume of patients with antrochoanal polyps, a condition that reduces nasal airflow without any causing anatomic changes (26). A recent study comparing oral breathing and nasal breathing showed the volume of the maxillary sinus to be smaller in patients who breathe through their mouth (27). This suggests that the factor impacting a change in the paranasal sinuses must have occurred in the developmental stage of the sinuses. While Koppe et al. (28) report to have found that an untreated cleft palate did not affect the maxillary sinus volume in adult patients, we found that an angulated hard palate did not affect the volume of the maxillary sinuses. This suggests that the key factor affecting the volume of the maxillary sinuses is the deviation of the nasal septum.

Previous studies that have calculated paranasal sinus volumes using several methods report that the calculation method had neither any influence on the results nor showed statistically significant differences between the results (7, 22, 29). We used the ImageJ software with the planimetry method and Cavalieri's principle in our calculations. While previous studies (6, 7, 21, 30) report maxillary sinus volumes to range from 11.1 cm³ to 23.0 cm³, our results were found to be consistent with those reported in the literature.

Conclusion

In the present study, we found that the presence of a septal deviation leads to reduced maxillary sinus volume on the deviated side regardless of whether or not it affects the os palatinum but does not affect the total volume. Results of previous studies and the present study suggest that nasal septal deviation has a significant impact on the development of the paranasal sinuses.

Ethics Committee Approval: Ethics committee approval was received for this study from the Ethics Committee of Firat University School of Medicine (2014-06/10).

Informed Consent: Informed consent was not received due to the retrospective nature of the study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - E.S.; Design - E.S.; Supervision - E.S., A.K.; Data Collection and/or Processing - E.S.; Analysis and/or Interpretation - E.S.; Literature Search - H.I.S.; Writing - E.S.; Critical Reviews - A.K., M.Ö.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Bu çalışma için etik kurul onayı Firat Üniversitesi Tıp Fakültesi Etik Kurulu'ndan alınmıştır (2014-06/10).

Hasta Onamı: Çalışmamızın geriye dönük tasarımından dolayı hasta onamı alınmamıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - E.S.; Tasarım - E.S.; Denetleme - E.S., A.K.; Veri Toplanması ve/veya İşlemesi - E.S.; Analiz ve/veya Yorum - E.S.; Literatür Taraması - H.I.S.; Yazıyı Yazan - E.S.; Eleştirel İnceleme - A.K., M.Ö.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

- Shapiro R, Schorr S. A consideration of the systemic factors that influence frontal sinus pneumatization. *Invest Radiol* 1980; 15: 191-202. [CrossRef]
- Fatu C, Puisoru M, Rotaru M, Truta AM. Morphometric evaluation of the frontal sinus in relation to age. *Ann Anat* 2006; 188: 275-80. [CrossRef]
- Cho SH, Kim TH, Kim KR, Lee JM, Lee DK, Kim JH, et al. Factors for maxillary sinus volume and craniofacial anatomical features in adults with chronic rhinosinusitis. *Arch Otolaryngol Head Neck Surg* 2010; 136: 610-5. [CrossRef]
- D'Ascanio L, Lancione C, Pompa G, Rebuffini E, Mansi N, Manzini M. Craniofacial growth in children with nasal septum deviation: a cephalometric comparative study. *Int J Pediatr Otorhinolaryngol* 2010; 74: 1180-3. [CrossRef]
- Sooknundun M, Kacker SK, Bhatia R, Deka RC. Nasal septal deviation: effective intervention and long term follow-up. *Int J Pediatr Otorhinolaryngol* 1986; 12: 65-72. [CrossRef]
- Kapusuz Gencer Z, Ozkiris M, Okur A, Karacavus S, Saydam L. The effect of nasal septal deviation on maxillary sinus volumes and development of maxillary sinusitis. *Eur Arch Otorhinolaryngol* 2013; 270: 3069-73. [CrossRef]
- Orhan I, Ormeci T, Aydin S, Altin G, Urger E, Soylu E, et al. Morphometric analysis of the maxillary sinus in patients with nasal septum deviation. *Eur Arch Otorhinolaryngol* 2014; 271: 727-32. [CrossRef]
- Park IH, Song JS, Choi H, Kim TH, Hoon S, Lee SH, et al. Volumetric study in the development of paranasal sinuses by CT imaging in Asian: a pilot study. *Int J Pediatr Otorhinolaryngol*. 2010; 74: 1347-50. [CrossRef]

9. Emami AJ, Brodsky L, Pizzuto M. Neonatal septoplasty: Case report and review of the literature. *Int J Pediatr Otorhinolaryngo*. 1996; 35: 271-5. [\[CrossRef\]](#)
10. Chintapatla S, Kudva YC, Nayar RC, Raghuvver TS, Prasad D. Septal deviation in neonates. *Indian Pediatr* 1989; 26: 678-82.
11. Korantzis A, Cardamakis E, Chelidonis E, Papamihalis T. Nasal septum deformity in the newborn infant during labour. *Eur J Obstet Gynecol Reprod Biol* 1992; 44: 41-6. [\[CrossRef\]](#)
12. Delaire J, Precious D. Influence of the nasal septum on maxillonasal growth in patients with congenital labiomaxillary cleft. *Cleft Palate J* 1986; 23: 270-7.
13. Holton NE, Yokley TR, Figueroa A. Nasal septal and craniofacial form in European- and African-derived populations. *J Anat* 2012; 221: 263-74. [\[CrossRef\]](#)
14. Grymer LF, Pallisgaard C, Melsen B. The nasal septum in relation to the development of the nasomaxillary complex: a study in identical twins. *Laryngoscope* 1991; 101: 863-8. [\[CrossRef\]](#)
15. Gray LP. The development and significance of septal and dental deformity from birth to eight years. *Int J Pediatr Otorhinolaryngol* 1983; 6: 265-77. [\[CrossRef\]](#)
16. Mladina R, Cujic E, Subaric M, Vukovic K. Nasal septal deformities in ear, nose, and throat patients: an international study. *Am J Otolaryngol* 2008; 29: 75-82. [\[CrossRef\]](#)
17. Klein JC. Nasal respiratory function and craniofacial growth. *Arch Otolaryngol Head Neck Surg* 1986; 112: 843-9. [\[CrossRef\]](#)
18. Parsons DS, Wald ER. Otitis media and sinusitis: similar diseases. *Otolaryngol Clin North Am* 1996; 29: 11-25.
19. Schlenker WL, Jennings BD, Jeiroudi MT, Caruso JM. The effects of chronic absence of active nasal respiration on the growth of the skull: a pilot study. *Am J Orthod Dentofacial Orthop* 2000; 117: 706-13. [\[CrossRef\]](#)
20. Jun BC, Song SW, Park CS, Lee DH, Cho KJ, Cho JH. The analysis of maxillary sinus aeration according to aging process; volume assessment by 3-dimensional reconstruction by high-resolution CT scanning. *Otolaryngol Head Neck Surg* 2005; 132: 429-34. [\[CrossRef\]](#)
21. Karakas S, Kavakli A. Morphometric examination of the paranasal sinuses and mastoid air cells using computed tomography. *Ann Saudi Med* 2005; 25: 41-5. [\[CrossRef\]](#)
22. Uchida Y, Goto M, Katsuki T, Akiyoshi T. A cadaveric study of maxillary sinus size as an aid in bone grafting of the maxillary sinus floor. *J Oral Maxillofac Surg* 1998; 56: 1158-63. [\[CrossRef\]](#)
23. Cho SH, Kim TH, Kim KR, Lee JM, Lee DK, Kim JH, et al. Factors for maxillary sinus volume and craniofacial anatomical features in adults with chronic rhinosinusitis. *Arch Otolaryngol Head Neck Surg* 2010; 136: 610-5. [\[CrossRef\]](#)
24. Barghouth G, Prior JO, Lepori D, Duvoisin B, Schnyder P, Gudinchet F. Paranasal sinuses in children: size evaluation of maxillary, sphenoid, and frontal sinuses by magnetic resonance imaging and proposal of volume index percentile curves. *Eur Radiol* 2002; 12: 1451-8. [\[CrossRef\]](#)
25. Firat AK, Miman MC, Firat Y, Karakas HM, Ozturan O, Altinok T. Effect of nasal septal deviation on total ethmoid cell volume. *J Laryngol Otol* 2006; 120: 200-4. [\[CrossRef\]](#)
26. Aydin S, Taskin U, Orhan I, Altas B, Oktay MF, Toksoz M, et al. The analysis of the maxillary sinus volumes and the nasal septal deviation in patients with antrochoanal polyps. *Eur Arch Otorhinolaryngol* 2015; 272: 3347-52. [\[CrossRef\]](#)
27. Agacayak KS, Gulsun B, Koparal M, Atalay Y, Aksoy O, Adiguzel O. Alterations in maxillary sinus volume among oral and nasal breathers. *Med Sci Monit* 2015; 21: 18-26. [\[CrossRef\]](#)
28. Koppe T, Weigel C, Barenklau M, Kaduk W, Bayerlein T, Gedrange T. Maxillary sinus pneumatization of an adult skull with an untreated bilateral cleft palate. *J Craniofac Surg* 2006; 34: 91-5. [\[CrossRef\]](#)
29. Degermenci M, Ertekin T, Ulger H, Acer N, Coskun A. The age-related development of maxillary sinus in children. *J Craniofac Surg* 2016; 27: 38-44. [\[CrossRef\]](#)
30. Pirner S, Tingelhoff K, Wagner I, Westphal R, Rilk M, Wahl FM, et al. CT-based manual segmentation and evaluation of paranasal sinuses. *Eur Arch Otorhinolaryngol* 2009; 266: 507-18. [\[CrossRef\]](#)