

identify the causes of airway narrowing that may clinically mimic adenoid hypertrophy. Evaluating a patient solely through radiological imaging such as lateral radiography may lead to overlooking benign or, in rare cases, malignant nasopharyngeal pathologies (3). Nasal endoscopy also allows for the assessment of the anatomical extent of the adenoid tissue prior to performing surgery, which may be clinically important given that the distribution of hypertrophy could affect symptomatology (4).

Endoscopic evaluation is also crucial for detecting rare vascular anomalies, such as an aberrant internal carotid artery (ICA). Unrecognized nasopharyngeal ICA aberrancy may lead to life-threatening hemorrhage during adenoidectomy. Nasal endoscopy may show a submucosal pulsatile mass in such cases (5). These considerations underscore the importance of thorough preoperative examination by otorhinolaryngologists before surgery.

I congratulate the authors for their contribution and believe these considerations may further strengthen the message of their study.

#### Footnotes

**Financial Disclosure:** The author declare that this study has received no financial support.

#### References

1. Hosri J, Hosn OA, Ghanem A, Daou AM, Ghadieh J, Zalaquett N, et al. Comparing flexible nasal endoscopy and lateral neck radiography when diagnosing children with adenoid hypertrophy: a case-control study. *Turk Arch Otorhinolaryngol.* 2025; 63: 185-9. [Crossref]
2. Pisutsiri N, Vathanophas V, Boonyabut P, Tritrakarn S, Vitayaudom N, Tanphaichitr A, et al. Adenoid measurement accuracy: a comparison of lateral skull film, flexible endoscopy, and intraoperative rigid endoscopy (gold standard). *Auris Nasus Larynx.* 2022; 49: 222-8. [Crossref]
3. Duarte VM, Liu YF, Shapiro NL. Uncommon presentation of a benign nasopharyngeal mass in an adolescent: comprehensive review of pediatric nasopharyngeal masses. *Case Rep Pediatr.* 2013; 2013: 816409. [Crossref]
4. Regmi D, Mathur NN, Bhattarai M. Rigid endoscopic evaluation of conventional curettage adenoidectomy. *J Laryngol Otol.* 2011; 125: 53-8. [Crossref]
5. Roje Ž, Kljajić Z, Kolić K, Ilić D. A surgical hazard during adenoidectomy in children-a case report. *Children (Basel).* 2025; 12: 139. [Crossref]

#### Reply

##### Dear Editor,

We would like to thank the author for their interest in our article entitled “Comparing Flexible Nasal Endoscopy and Lateral Neck Radiography When Diagnosing Children with Adenoid Hypertrophy” and for their thoughtful and constructive comments.

We appreciate the emphasis on several important aspects of adenoid evaluation in pediatric patients. In particular, the points raised regarding the limitations of lateral neck radiography, including radiation exposure and potential variability during image acquisition, are well taken. These factors are indeed relevant in daily clinical practice and should be considered when selecting diagnostic modalities.

We also agree that flexible nasal endoscopy offers significant advantages by enabling direct visualization of the nasopharynx. As highlighted, endoscopic assessment may provide additional diagnostic information beyond the estimation of adenoid size, including the identification of alternative causes of nasopharyngeal obstruction. Furthermore, the role of endoscopy in preoperative evaluation, particularly in recognizing rare but potentially critical conditions such as vascular anomalies, is of clear clinical importance.

At the same time, we would like to emphasize that lateral neck radiography may still have a role in selected clinical settings. In situations where endoscopic examination is not feasible, limited, or not well tolerated, radiography can serve as a useful adjunctive tool. Therefore, rather than considering these modalities as mutually exclusive, they may be better viewed as complementary, depending on the clinical context and available resources.

Overall, we believe that the author’s comments enrich the discussion and highlight important diagnostic and safety considerations in the evaluation of adenoid hypertrophy.

We thank the author again for their valuable contribution.

Sincerely,

✉ Jad Hosri<sup>1</sup>, ✉ Omar Aboul Hosn<sup>1</sup>, ✉ Anthony Ghanem<sup>1</sup>,  
✉ Anne Marie Daou<sup>1</sup>, ✉ Justin Ghadieh<sup>1</sup>, ✉ Nader Zalaquett<sup>2</sup>,  
✉ Randa Barazi<sup>1</sup>

<sup>1</sup>American University of Beirut Medical Center, Department of Otolaryngology-Head and Neck Surgery, Beirut, Lebanon

<sup>2</sup>American University of Beirut Faculty of Medicine, Beirut, Lebanon